

Introduction to prismatic cohomology

June 20, 2023

Note: generally everything is sourced from Bhatt–Scholze [2] unless otherwise specified.

Motivation: perfectoid rings

(Integral) perfectoid rings:

- Analogue of perfect rings for \mathbb{Z}_p -algebras

Motivation: perfectoid rings

(Integral) perfectoid rings:

- Analogue of perfect rings for \mathbb{Z}_p -algebras
- p -complete \mathbb{Z}_p -algebras such that mod p , Frobenius is a bijection; \exists p th power dividing p ; some other technical conditions

Motivation: perfectoid rings

(Integral) perfectoid rings:

- Analogue of perfect rings for \mathbb{Z}_p -algebras
- p -complete \mathbb{Z}_p -algebras such that mod p , Frobenius is a bijection; \exists p th power dividing p ; some other technical conditions
- Examples: perfect \mathbb{F}_p -algebras, $\mathbb{Z}_p[p^{1/p^\infty}]_p^\wedge$

Motivation: perfectoid rings

(Integral) perfectoid rings:

- Analogue of perfect rings for \mathbb{Z}_p -algebras
- p -complete \mathbb{Z}_p -algebras such that mod p , Frobenius is a bijection; $\exists p$ th power dividing p ; some other technical conditions
- Examples: perfect \mathbb{F}_p -algebras, $\mathbb{Z}_p[p^{1/p^\infty}]_p^\wedge$
- Main interest: tilting correspondence: for R perfectoid, $\{\text{perfectoid } R\text{-algebras}\} \leftrightarrow \{\text{perfectoid } R^b\text{-algebras}\}$ (where $R^b = \lim_{\phi} R/p$)

Motivation: perfectoid rings

(Integral) perfectoid rings:

- Analogue of perfect rings for \mathbb{Z}_p -algebras
- p -complete \mathbb{Z}_p -algebras such that mod p , Frobenius is a bijection; $\exists p$ th power dividing p ; some other technical conditions
- Examples: perfect \mathbb{F}_p -algebras, $\mathbb{Z}_p[p^{1/p^\infty}]_p^\wedge$
- Main interest: tilting correspondence: for R perfectoid, $\{\text{perfectoid } R\text{-algebras}\} \leftrightarrow \{\text{perfectoid } R^b\text{-algebras}\}$ (where $R^b = \lim_{\phi} R/p$)
- Example: Fontaine–Winterberger theorem

$$\mathrm{Gal}_{\mathbb{Q}_p(p^{1/p^\infty})_p^\wedge} \simeq \mathrm{Gal}_{\mathbb{F}_p((T^{1/p^\infty}))_T^\wedge}$$

Obstructions

Perfectoid condition is unpleasantly restrictive. Possible solutions:

Obstructions

Perfectoid condition is unpleasantly restrictive. Possible solutions:

- Take covers by perfectoid spaces and tilt keeping track of the covers (leading to theory of diamonds)

Obstructions

Perfectoid condition is unpleasantly restrictive. Possible solutions:

- Take covers by perfectoid spaces and tilt keeping track of the covers (leading to theory of diamonds)
- Generalize: if perfectoid spaces are analogous to perfect \mathbb{F}_p -algebras, how can we drop the “perfect”?

Perfect prisms

Let's reformulate perfectoid rings using the A_{inf} functor

$$A_{\text{inf}}(R) = W(R^b).$$

Perfect prisms

Let's reformulate perfectoid rings using the A_{inf} functor

$$A_{\text{inf}}(R) = W(R^b).$$

Theorem

There exists a surjection $\theta : A_{\text{inf}}(R) \rightarrow R$ with kernel (ξ) principal.

Perfect prisms

Let's reformulate perfectoid rings using the A_{inf} functor

$$A_{\text{inf}}(R) = W(R^b).$$

Theorem

There exists a surjection $\theta : A_{\text{inf}}(R) \rightarrow R$ with kernel (ξ) principal.

So the data of $A_{\text{inf}}(R)$ (depending only on R^b) and (ξ) together recover R .

Perfect prisms

What’s the codomain of this functor?

Perfect prisms

What’s the codomain of this functor?

- $A_{\text{inf}}(R)$ is equipped with a lift of Frobenius ϕ (or better, δ -structure)
- (ξ) is a principal ideal with “distinguished generator” ($\delta(\xi)$ is a unit, or equivalently $p \in (\xi, \phi(\xi))$), such that $A_{\text{inf}}(R)$ is (derived) (p, ξ) -complete.

Perfect prisms

What’s the codomain of this functor?

- $A_{\text{inf}}(R)$ is equipped with a lift of Frobenius ϕ (or better, δ -structure)
- (ξ) is a principal ideal with “distinguished generator” ($\delta(\xi)$ is a unit, or equivalently $p \in (\xi, \phi(\xi))$), such that $A_{\text{inf}}(R)$ is (derived) (p, ξ) -complete.

Further since R^b is perfect, ϕ is an isomorphism.

Definition

A **perfect prism** is a pair (A, I) where A is a δ -ring such that the Frobenius is an isomorphism and I is a principal ideal such that $p \in (I, \phi(I))$ and A is derived (p, I) -complete.

Perfect prisms

Theorem

The functors $R \mapsto (A_{\text{inf}}(R), \ker(A_{\text{inf}}(R) \rightarrow R))$ and $(A, I) \mapsto A/I$ are inverse functors defining an equivalence between perfectoid rings and perfect prisms.

Perfect prisms

Theorem

The functors $R \mapsto (A_{\text{inf}}(R), \ker(A_{\text{inf}}(R) \rightarrow R))$ and $(A, I) \mapsto A/I$ are inverse functors defining an equivalence between perfectoid rings and perfect prisms.

Example

$\mathbb{F}_p = \mathbb{Z}_p/(p) \mapsto (\mathbb{Z}_p, (p))$; more generally any perfect \mathbb{F}_p -algebra R corresponds to $(W(R), (p))$.

Perfect prisms

Theorem

The functors $R \mapsto (A_{\text{inf}}(R), \ker(A_{\text{inf}}(R) \rightarrow R))$ and $(A, I) \mapsto A/I$ are inverse functors defining an equivalence between perfectoid rings and perfect prisms.

Example

$\mathbb{F}_p = \mathbb{Z}_p/(p) \mapsto (\mathbb{Z}_p, (p))$; more generally any perfect \mathbb{F}_p -algebra R corresponds to $(W(R), (p))$.

A mixed-characteristic example is

$$\mathbb{Z}_p[p^{1/p^\infty}]_p^\wedge \mapsto (\mathbb{Z}_p[[T^{1/p^\infty}]]_p^\wedge, (T - p)).$$

Perfect prisms

Theorem

The functors $R \mapsto (A_{\text{inf}}(R), \ker(A_{\text{inf}}(R) \rightarrow R))$ and $(A, I) \mapsto A/I$ are inverse functors defining an equivalence between perfectoid rings and perfect prisms.

Example

$\mathbb{F}_p = \mathbb{Z}_p/(p) \mapsto (\mathbb{Z}_p, (p))$; more generally any perfect \mathbb{F}_p -algebra R corresponds to $(W(R), (p))$.

A mixed-characteristic example is

$$\mathbb{Z}_p[p^{1/p^\infty}]_p^\wedge \mapsto (\mathbb{Z}_p[[T^{1/p^\infty}]]_p^\wedge, (T - p)).$$

To “deperfect” perfectoid rings, we should look for a version without the “perfect” condition.

Prisms

The “perfectness” is the condition that ϕ be an isomorphism, and it’s geometrically better to ask for I locally principal. So:

Prisms

The “perfectness” is the condition that ϕ be an isomorphism, and it’s geometrically better to ask for I locally principal. So:

Definition

A **prism** is a pair (A, I) where A is any δ -ring and I is a locally principal ideal such that $p \in (I, \phi(I))$ and A is (p, I) -complete.

A morphism of prisms is a map of δ -rings compatible with the chosen ideals.

Prisms

Prisms can be:

- **perfect** if ϕ is an isomorphism (as above);

Prisms

Prisms can be:

- **perfect** if ϕ is an isomorphism (as above);
- **bounded** if A/I has bounded p^∞ -torsion (technical condition);

Prisms

Prisms can be:

- **perfect** if ϕ is an isomorphism (as above);
- **bounded** if A/I has bounded p^∞ -torsion (technical condition);
- **orientable** if I is principal, or **oriented** if a generator (orientation) is given (can usually be assumed for convenience);

Prisms

Prisms can be:

- **perfect** if ϕ is an isomorphism (as above);
- **bounded** if A/I has bounded p^∞ -torsion (technical condition);
- **orientable** if I is principal, or **oriented** if a generator (orientation) is given (can usually be assumed for convenience);
- **crystalline** if $I = (p)$ (special case).

Prisms

Prisms can be:

- **perfect** if ϕ is an isomorphism (as above);
- **bounded** if A/I has bounded p^∞ -torsion (technical condition);
- **orientable** if I is principal, or **oriented** if a generator (orientation) is given (can usually be assumed for convenience);
- **crystalline** if $I = (p)$ (special case).

It turns out $(A, (p))$ is a prism iff A is p -torsion-free.

Tilting equivalence

We can recover the tilting equivalence from a few lemmas about prisms:

Tilting equivalence

We can recover the tilting equivalence from a few lemmas about prisms:

Lemma (Rigidity)

If $(A, I) \rightarrow (B, J)$ is a map of prisms, then $J = IB$.

Tilting equivalence

We can recover the tilting equivalence from a few lemmas about prisms:

Lemma (Rigidity)

If $(A, I) \rightarrow (B, J)$ is a map of prisms, then $J = IB$.

Lemma (Lifting)

If (A, I) is a perfect prism and (B, J) is any prism, then any map $A/I \rightarrow B/J$ lifts uniquely to a prism map $(A, I) \rightarrow (B, J)$.

Tilting equivalence

We can recover the tilting equivalence from a few lemmas about prisms:

Lemma (Rigidity)

If $(A, I) \rightarrow (B, J)$ is a map of prisms, then $J = IB$.

Lemma (Lifting)

If (A, I) is a perfect prism and (B, J) is any prism, then any map $A/I \rightarrow B/J$ lifts uniquely to a prism map $(A, I) \rightarrow (B, J)$.

Corollary

Let R be a perfectoid ring with tilt R^{\flat} . Then the categories of perfectoid R -algebras and perfectoid R^{\flat} -algebras are equivalent.

Tilting equivalence

Corollary

Let R be a perfectoid ring with tilt R^{\flat} . Then the categories of perfectoid R -algebras and perfectoid R^{\flat} -algebras are equivalent.

Proof.

Write R uniquely as A/I for a perfect prism (A, I) . Then maps $R \rightarrow R'$ lift uniquely to $(A, I) \rightarrow (A', I')$ and so induce $A/p = R^{\flat} \rightarrow A'/p = R'^{\flat}$ and similarly in reverse, which gives a bijection by rigidity. □

Tilting equivalence

Corollary

Let R be a perfectoid ring with tilt R^{\flat} . Then the categories of perfectoid R -algebras and perfectoid R^{\flat} -algebras are equivalent.

Proof.

Write R uniquely as A/I for a perfect prism (A, I) . Then maps $R \rightarrow R'$ lift uniquely to $(A, I) \rightarrow (A', I')$ and so induce $A/p = R^{\flat} \rightarrow A'/p = R'^{\flat}$ and similarly in reverse, which gives a bijection by rigidity. □

In fact we have no restriction on (A', I') so actually this is a stronger version!

Prismatic site

For the tilting equivalence we study perfectoid algebras over a perfectoid R , or equivalently perfect prisms (A, I) together with a map $R \rightarrow A/I$. Now we want to generalize to non-perfectoid rings:

Definition

The **(absolute) prismatic site** $(R)_{\Delta}$ of a ring R is the category of prisms (A, I) together with a map $R \rightarrow A/I$, equipped with the $[BZZZT]$ topology.

Prismatic site

For the tilting equivalence we study perfectoid algebras over a perfectoid R , or equivalently perfect prisms (A, I) together with a map $R \rightarrow A/I$. Now we want to generalize to non-perfectoid rings:

Definition

The **(absolute) prismatic site** $(R)_{\Delta}$ of a ring R is the category of prisms (A, I) together with a map $R \rightarrow A/I$, equipped with the $[BZZZT]$ topology.

There is also a relative version, which is what we'll mostly use:

Definition

For a fixed prism (A, I) and an A/I -algebra R , the **prismatic site** $(R/A)_{\Delta}$ of R over (A, I) is the category of prisms (B, J) together with compatible maps $(A, I) \rightarrow (B, J)$ and $R \rightarrow B/J$, equipped with the $[BZZZT]$ topology.

Aside

Note: these are not really sites, because the arrows are the wrong way! Really these should be the opposites of these categories, which works well for replacing R by a scheme (e.g. $\text{Spec } A/I \rightarrow \text{Spec } R$).

Sheaves and cohomology

There are several natural sheaves on the prismatic sites:

Sheaves and cohomology

There are several natural sheaves on the prismatic sites:

- $\mathcal{O}_{\Delta} : (A, I) \mapsto A$ (the structure sheaf)

Sheaves and cohomology

There are several natural sheaves on the prismatic sites:

- $\mathcal{O}_{\Delta} : (A, I) \mapsto A$ (the structure sheaf)
- $\overline{\mathcal{O}}_{\Delta} : (A, I) \mapsto A/I$ (the Hodge–Tate sheaf)

Sheaves and cohomology

There are several natural sheaves on the prismatic sites:

- $\mathcal{O}_{\Delta} : (A, I) \mapsto A$ (the structure sheaf)
- $\overline{\mathcal{O}}_{\Delta} : (A, I) \mapsto A/I$ (the Hodge–Tate sheaf)
- $\mathcal{I}_{\Delta} : (A, I) \mapsto I$ (the Hodge–Tate ideal sheaf)

Sheaves and cohomology

There are several natural sheaves on the prismatic sites:

- $\mathcal{O}_{\Delta} : (A, I) \mapsto A$ (the structure sheaf)
- $\overline{\mathcal{O}}_{\Delta} : (A, I) \mapsto A/I$ (the Hodge–Tate sheaf)
- $\mathcal{I}_{\Delta} : (A, I) \mapsto I$ (the Hodge–Tate ideal sheaf)

all of which can be defined on either the absolute or relative prismatic sites.

Sheaves and cohomology

There are several natural sheaves on the prismatic sites:

- $\mathcal{O}_{\Delta} : (A, I) \mapsto A$ (the structure sheaf)
- $\overline{\mathcal{O}}_{\Delta} : (A, I) \mapsto A/I$ (the Hodge–Tate sheaf)
- $\mathcal{I}_{\Delta} : (A, I) \mapsto I$ (the Hodge–Tate ideal sheaf)

all of which can be defined on either the absolute or relative prismatic sites.

Taking cohomology gives

- $\Delta_{R/A} = R\Gamma((R/A)_{\Delta}, \mathcal{O}_{\Delta})$, the relative prismatic cohomology;

Sheaves and cohomology

There are several natural sheaves on the prismatic sites:

- $\mathcal{O}_\Delta : (A, I) \mapsto A$ (the structure sheaf)
- $\overline{\mathcal{O}}_\Delta : (A, I) \mapsto A/I$ (the Hodge–Tate sheaf)
- $\mathcal{I}_\Delta : (A, I) \mapsto I$ (the Hodge–Tate ideal sheaf)

all of which can be defined on either the absolute or relative prismatic sites.

Taking cohomology gives

- $\Delta_{R/A} = R\Gamma((R/A)_\Delta, \mathcal{O}_\Delta)$, the relative prismatic cohomology;
- $\Delta_R = R\Gamma((R)_\Delta, \mathcal{O}_\Delta)$, the absolute prismatic cohomology;

Sheaves and cohomology

There are several natural sheaves on the prismatic sites:

- $\mathcal{O}_{\Delta} : (A, I) \mapsto A$ (the structure sheaf)
- $\overline{\mathcal{O}}_{\Delta} : (A, I) \mapsto A/I$ (the Hodge–Tate sheaf)
- $\mathcal{I}_{\Delta} : (A, I) \mapsto I$ (the Hodge–Tate ideal sheaf)

all of which can be defined on either the absolute or relative prismatic sites.

Taking cohomology gives

- $\Delta_{R/A} = R\Gamma((R/A)_{\Delta}, \mathcal{O}_{\Delta})$, the relative prismatic cohomology;
- $\Delta_R = R\Gamma((R)_{\Delta}, \mathcal{O}_{\Delta})$, the absolute prismatic cohomology;
- $\overline{\Delta}_{R/A} = R\Gamma((R/A)_{\Delta}, \overline{\mathcal{O}}_{\Delta})$, relative Hodge–Tate cohomology, and similarly for the absolute version

Prismatic cohomology

At least in the relative case,

$$\overline{\Delta}_{R/A} = \Delta_{R/A} \otimes_A A/I$$

so prismatic cohomology carries the most information.

Prismatic cohomology

At least in the relative case,

$$\overline{\Delta}_{R/A} = \Delta_{R/A} \otimes_A A/I$$

so prismatic cohomology carries the most information.

Example

Suppose $R = A/I$. Then

$$\Delta_{R/A} = A, \quad \overline{\Delta}_{R/A} = A/I = R.$$

Perfectoidization

We're familiar with perfection of \mathbb{F}_p -algebras: $R_{\text{perf}} = \text{colim}_{\phi} R$.
The analogue for mixed-characteristic rings is **perfectoidization**.

Perfectoidization

We're familiar with perfection of \mathbb{F}_p -algebras: $R_{\text{perf}} = \text{colim}_{\phi} R$.
The analogue for mixed-characteristic rings is **perfectoidization**.
First, there's a straightforward analogy for δ -rings:
 $A_{\text{perf}} = (\text{colim}_{\phi} A)^{\wedge}$.

Perfectoidization

We're familiar with perfection of \mathbb{F}_p -algebras: $R_{\text{perf}} = \text{colim}_{\phi} R$.
The analogue for mixed-characteristic rings is **perfectoidization**.

First, there's a straightforward analogy for δ -rings:

$$A_{\text{perf}} = (\text{colim}_{\phi} A)^{\wedge}.$$

Then we can define the perfectoidization in two ways: first, let R be an S -algebra for S perfectoid, so we can write $S = A/I$ for a perfect prism (A, I) . Then

$$R_{\text{perfd}} := \Delta_{R/A, \text{perf}} \otimes_A^{\mathbb{L}} A/I.$$

Perfectoidization

This is sort of a “perfected Hodge–Tate cohomology.” Note that it is in general a derived ring (in your preferred model, e.g. an animated ring or a commutative algebra object in $D(R)$). However a priori it depends on the choice of a presentation $A/I = S \rightarrow R$.

Perfectoidization

This is sort of a “perfected Hodge–Tate cohomology.” Note that it is in general a derived ring (in your preferred model, e.g. an animated ring or a commutative algebra object in $D(R)$). However a priori it depends on the choice of a presentation $A/I = S \rightarrow R$.

Fortunately we can define this in another way which makes the independence clear by modifying the site: let $(R)_{\Delta}^{\text{perf}}$ be the **(absolute) perfect prismatic site** of R , consisting of **perfect** prisms (A, I) with maps $R \rightarrow A/I$. Then

$$R_{\text{perfd}} \simeq R\Gamma((R)_{\Delta}^{\text{perf}}, \overline{\mathcal{O}}_{\Delta}).$$

Comparison theorems

The other big motivation for prismatic cohomology is to generalize and unify various classical p -adic cohomology theories. For example:

Comparison theorems

The other big motivation for prismatic cohomology is to generalize and unify various classical p -adic cohomology theories. For example:

- Crystalline cohomology

Comparison theorems

The other big motivation for prismatic cohomology is to generalize and unify various classical p -adic cohomology theories. For example:

- Crystalline cohomology
- Hodge cohomology

Comparison theorems

The other big motivation for prismatic cohomology is to generalize and unify various classical p -adic cohomology theories. For example:

- Crystalline cohomology
- Hodge cohomology
- de Rham cohomology


Comparison theorems

The other big motivation for prismatic cohomology is to generalize and unify various classical p -adic cohomology theories. For example:

- Crystalline cohomology
- Hodge cohomology
- de Rham cohomology
- p -adic étale cohomology of the generic fiber


Comparison theorems

This is the origin of the picture motivating the term "prism"
 (stolen from [1, Lecture 1]) over $\text{Spec } \mathbb{Z}_p[[u]]$ (should really be Σ):



Comparison theorems

This is the origin of the picture motivating the term "prism"
 (stolen from [1, Lecture 1]) over $\text{Spec } \mathbb{Z}_p[[u]]$ (should really be Σ):



We'll leave details for future talks, but let's spell out the comparison in one example:

Crystalline comparison

Suppose (A, I) is crystalline and R is an A/I -algebra. Then $\Delta_{R/A}$ is **almost** equal to the classical crystalline cohomology $R\Gamma_{\text{crys}}(R/A)$:

$$R\Gamma_{\text{crys}}(R/A) = \phi^* \Delta_{R/A} = \Delta_{R/A} \widehat{\otimes}_{A, \phi_A}^{\mathbb{L}} A.$$

Crystalline comparison

Suppose (A, I) is crystalline and R is an A/I -algebra. Then $\Delta_{R/A}$ is **almost** equal to the classical crystalline cohomology $R\Gamma_{\text{crys}}(R/A)$:

$$R\Gamma_{\text{crys}}(R/A) = \phi^* \Delta_{R/A} = \Delta_{R/A} \widehat{\otimes}_{A, \phi_A}^{\mathbb{L}} A.$$

When (A, I) is any perfect prism, we can then think of $\Delta_{R/A}$ as generalizing crystalline cohomology to mixed-characteristic perfectoids (after correcting by a Frobenius twist).

Qrsp rings and the Nygaard filtration

There is a site of \mathbb{Z}_p -algebras called the quasisyntomic site, with a basis of quasiregular semiperfectoid (qrsp) rings (roughly, those which are quotients of perfectoid rings by quasiregular ideals). Prismatic cohomology is well-behaved on these:

Proposition

If R is qrsp, the absolute prismatic site $(R)_{\Delta}$ has an initial object $(\Delta_R^{\text{init}}, (d))$, and for any prism (A, I) with a map $A/I \rightarrow R$ we have

$$\Delta_{R/A} = \Delta_R = \Delta_R^{\text{init}}.$$

In particular the prismatic cohomology of R is concentrated in degree 0, and $R\Gamma((R)_{\Delta}, \mathcal{I}_{\Delta}) = d\Delta_R$ is principal.

Qrsp rings and the Nygaard filtration

Thus we can define a filtration $\mathcal{N}^{\geq i} \Delta_R = \phi^{-1}(d^i \Delta_R)$, or (equivalently) $\mathcal{N}^{\geq i} \Delta_{R/A} = \phi^{-1}(d^i \Delta_{R/A})$.

Qrsp rings and the Nygaard filtration

Thus we can define a filtration $\mathcal{N}^{\geq i} \Delta_R = \phi^{-1}(d^i \Delta_R)$, or (equivalently) $\mathcal{N}^{\geq i} \Delta_{R/A} = \phi^{-1}(d^i \Delta_{R/A})$.

Then by descent we can define this filtration on the prismatic cohomology (absolute or relative) of any quasisyntomic ring R . Write $\mathcal{N}^i \Delta_R$, $\mathcal{N}^i \Delta_{R/A}$ for the graded pieces, and $\hat{\Delta}_R$, $\hat{\Delta}_{R/A}$ for the completion with respect to this filtration.

Topological connections

Now let's briefly review topological Hochschild homology: for an E_∞ -ring spectrum R (which for us will be a usual discrete ring), $\mathrm{THH}(R)$ can be defined as the universal S^1 -equivariant E_∞ -ring spectrum over R . The S^1 -action means we can form $\mathrm{TC}^-(R) := \mathrm{THH}(R)^{hS^1}$ and $\mathrm{TP}(R) := \mathrm{THH}(R)^{tS^1}$, which come with a natural map $\mathrm{TC}^-(R) \rightarrow \mathrm{TP}(R)$. (We take all of these with \mathbb{Z}_p -coefficients, i.e. p -completed.)

Topological connections

Now let's briefly review topological Hochschild homology: for an E_∞ -ring spectrum R (which for us will be a usual discrete ring), $\mathrm{THH}(R)$ can be defined as the universal S^1 -equivariant E_∞ -ring spectrum over R . The S^1 -action means we can form $\mathrm{TC}^-(R) := \mathrm{THH}(R)^{hS^1}$ and $\mathrm{TP}(R) := \mathrm{THH}(R)^{tS^1}$, which come with a natural map $\mathrm{TC}^-(R) \rightarrow \mathrm{TP}(R)$. (We take all of these with \mathbb{Z}_p -coefficients, i.e. p -completed.)

Bökstedt's theorem: $\pi_{2*} \mathrm{THH}(\mathbb{F}_p) = \mathbb{F}_p$, and odd homotopy groups vanish. There is a generalization to perfectoid rings R : $\pi_{2*} \mathrm{THH}(R) = R$, and odd groups vanish.

Topological connections

Now let's briefly review topological Hochschild homology: for an E_∞ -ring spectrum R (which for us will be a usual discrete ring), $\mathrm{THH}(R)$ can be defined as the universal S^1 -equivariant E_∞ -ring spectrum over R . The S^1 -action means we can form $\mathrm{TC}^-(R) := \mathrm{THH}(R)^{hS^1}$ and $\mathrm{TP}(R) := \mathrm{THH}(R)^{tS^1}$, which come with a natural map $\mathrm{TC}^-(R) \rightarrow \mathrm{TP}(R)$. (We take all of these with \mathbb{Z}_p -coefficients, i.e. p -completed.)

Bökstedt's theorem: $\pi_{2*} \mathrm{THH}(\mathbb{F}_p) = \mathbb{F}_p$, and odd homotopy groups vanish. There is a generalization to perfectoid rings R : $\pi_{2*} \mathrm{THH}(R) = R$, and odd groups vanish.

What about qrsp rings?

Topological connections

The evenness is still true, but the description is more complicated:
 $\pi_{2*} \mathrm{THH}(R) = \mathcal{N}^n \hat{\Delta}_R \{*\}$! (Here $\{*\}$ is a twist we'll skim over.)

Topological connections

The evenness is still true, but the description is more complicated:

$\pi_{2*} \mathrm{THH}(R) = \mathcal{N}^n \hat{\Delta}_R \{*\}$! (Here $\{*\}$ is a twist we’ll skim over.)

Moreover, $\mathrm{TC}^-(R)$ and $\mathrm{TP}(R)$ also have even homotopy groups, and

$$\pi_{2*} \mathrm{TC}^-(R) = \mathcal{N}^{\geq *}\hat{\Delta}_R \{*\}, \quad \pi_{2*} \mathrm{TP}(R) = \hat{\Delta}_R \{*\}.$$

Topological connections

The evenness is still true, but the description is more complicated:
 $\pi_{2*} \mathrm{THH}(R) = \mathcal{N}^n \hat{\Delta}_R \{*\}$! (Here $\{*\}$ is a twist we’ll skim over.)
 Moreover, $\mathrm{TC}^-(R)$ and $\mathrm{TP}(R)$ also have even homotopy groups,
 and

$$\pi_{2*} \mathrm{TC}^-(R) = \mathcal{N}^{\geq *}\hat{\Delta}_R \{*\}, \quad \pi_{2*} \mathrm{TP}(R) = \hat{\Delta}_R \{*\}.$$

In particular $\pi_0 \mathrm{TC}^-(R) = \pi_0 \mathrm{TP}(R) = \hat{\Delta}_R$. Since qrsp rings form a basis for the quasisyntomic site, it follows that the quasisyntomic sheafification of $\pi_0 \mathrm{TC}^-(-) = \pi_0 \mathrm{TP}(-)$ is $\hat{\Delta}_-$.

Topological connections

One can use the cyclotomic structure on THH to get Frobenius maps $TC^-(R) \rightarrow TP(R)$ and take the equalizer to get $TC(R)$; doing the same sheafification process gives the **syntomic cohomology** of R , which previously was not defined in this generality. One can also expand to even ring spectra other than $THH(R)$ for R qrsp to get the even and motivic filtrations [4].

Topological connections

One can use the cyclotomic structure on THH to get Frobenius maps $TC^-(R) \rightarrow TP(R)$ and take the equalizer to get $TC(R)$; doing the same sheafification process gives the **syntomic cohomology** of R , which previously was not defined in this generality. One can also expand to even ring spectra other than $THH(R)$ for R qrsp to get the even and motivic filtrations [4].

In particular, p -locally the cyclotomic trace $K(R) \rightarrow TC(R)$ is an equivalence by [3], so we can say something about K-theory as a consequence: e.g. quasisyntomic locally its homotopy groups are even, i.e. the quasisyntomic sheafification of $\pi_n K(-, \mathbb{Z}_p)$ vanishes for n odd.

References



Bhargav Bhatt.

Prismatic cohomology: {E}ilenberg lectures.

Lecture notes available at <http://www-personal.umich.edu/~bhattb/teaching/prismatic-columbia/>, year=2022.



Bhargav Bhatt and Peter Scholze.

Prisms and prismatic cohomology.

Annals of Mathematics, 196(3):1135–1275, 2022.



Dustin Clausen, Akhil Mathew, and Matthew Morrow.

K -theory and topological cyclic homology of henselian pairs.

Journal of the American Mathematical Society, 34(2):411–473, 2021.



Jeremy Hahn, Arpon Raksit, and Dylan Wilson.

A motivic filtration on the topological cyclic homology of commutative ring spectra.